

infestans

[1].

3

5×7×30

3-5

2-3

5-7

80-

6

P.infestans

P.

: 0 –

, 5 –

, 1-4 –

: $X = \sum (I_i \times D_i) / n$, I_i –

; D_i –

; n –

7-20

AAS-3,

[2].

40%

70-

[8],

[9].

()

(=0,0003)

(=0,0009).

				Fe	Cu
0	9,8±0,49	31,7±0,48	11,68±0,52	2,095±0,17	
24	10,4±0,45	40,5±1,95	15,2±0,77	2,236±0,25	
76	14,2±0,72	58,3±1,43	19,23±0,97	2,711±0,303	

[3-5].

(160

500

14

1 . [6].

()

(5).

48

1

0,06*

(R=0,98; F=27,6; p=0,035 (. 1),

= 31,87 + 0,349*

(R=0,99; F=3598; p=0,0053 (. 2).

[7].

(10

(p=0,0001),

(p=0,0014),

(p=0,004).


(p<0,0001),

p=0,015.


r=0,66, p=0,032,

r=0,82,

($p > 0,05$)



1.



2.

[10].

1. , 2003. 31 . 2. . // . 2005.
4. . 74-91. 3. Levine A., Tenhaken R.//Cell. 1994.V. 79. N 4. P. 583-593.
4. Mittler R., Herr E.H.//Proc. Natl. Acad. Sci. USA. 1999. V. 96. N 24. P. 14165-14170. 5. Mittler R., Feng X., Cohen M.//Plant Cell. 1998. V.10. N3. P. 461-473. 6. Phytophthora infestans// . – 1981. – 6.- . 470-472. 7. Filippov, A. V., Kuznetsova, M. A., Rogozhin, A. N., Shirokova, E. A. & Kostjashov, V. V. Systemic resistance of potatoes to late blight activated by preplanting treatment of tubers with specially formed low frequency pulse electric fields//J. Rus. Phytopathol. Soc. 2000. N 1. P. 69-73. 8. , 2003. 9. , 2006. 10. Phytophthora infestans (Mont.) de Bary