

• • • • • , • • • • • , • • • • • , • • • • • ,

1,0-1,8 %, - 2-4

5,0-5,5 /100

100<sup>2</sup> 25-50<sup>2</sup>

2007 .

0,25-0,27

0,25-0,27

0,60 [1].

75-80%

[2,3].

( )

[4,5].

, / ,						
(2007-2008 .)						
	2007	2008		2007	2008	
20,5	21,2	20,8	21,0	22,0	21,5	
22,8	23,6	23,2	23,7	24,7	24,2	
23,9	24,9	24,4	25,1	26,2	25,6	
27,4	28,5	28,0	28,9	30,5	29,7	

60%  
2-4

8-10°

2,42 / .

(1,2

1 )

2,72 / .

3,6 /

[5].

3

3 8 %.

1. , . . . . . 2 / . . . . . :- . :  
, 1965 2. , . . . . .  
/ . . . . . //  
, 1991 - N 10. 3. , . . . . .  
. . . . . // .- . 1991. 4.  
, . . . . . /  
. . . . . // . . . . . -  
. 1993 5. , . . . . .  
/ . . . . .  
// . - . . . . - , 2004.

- 176 .

**APPLICATION OF MIZORIN AND NATURAL BISHOFITE TO POTATO GROWN ON A DECOMPACTED SOIL IN THE  
NORTHERN CASPIAN REGION**

**A.Y. Moskvichev, A.V. Balashov, V.V. Pyatibratov** *Volgograd State Agricultural Academy, Universitetskii pr. 26, Volgograd, 400002  
Russia, e-mail: [VGSXA@avtlg.ru](mailto:VGSXA@avtlg.ru)*

**Summary.** *Problems related to the improvement of potato growing technology with the use of the biopreparation Mizorin and the natural mineral bishofite on a decompacted light-chestnut soil of the Northern Caspian region were considered*

**Key words:** *potato, Mizorin, bishofite, tuber yield, soil decompaction*